

Optical Identification of the Hardest X-ray Source in the *ASCA* Large Sky Survey

Masayuki Akiyama^{1,2,3} and Kouji Ohta^{1,2}

Department of Astronomy, Kyoto University, Kyoto 606-8502, Japan

Toru Yamada¹

Astronomical Institute, Tohoku University, Sendai 980-8578, Japan

Michael Eracleous^{1,4}

Department of Astronomy, University of California, Berkeley, CA 94720

Jules P. Halpern¹

Department of Astronomy, Columbia University, New York, NY 10027

Nobunari Kashikawa and Masafumi Yagi

National Astronomical Observatory, Mitaka, Tokyo 181-8588, Japan

Wataru Kawasaki

Department of Astronomy, University of Tokyo, Tokyo 113-8658, Japan

Masaaki Sakano³ and Takeshi Tsuru

Department of Physics, Kyoto University, Kyoto 606-8502, Japan

and

Yoshihiro Ueda and Tadayuki Takahashi

Institute of Space and Astronautical Science, Kanagawa 229-8510, Japan

ABSTRACT

We report the optical identification of the hardest X-ray source (AX J131501+3141) detected in an unbiased wide-area survey in the 0.5–10 keV band, the *ASCA* Large Sky Survey (LSS). The X-ray spectrum of the source is very hard and is well reproduced by a power law component ($\Gamma = 1.5^{+0.7}_{-0.6}$) with $N_{\text{H}} = 6^{+4}_{-2} \times 10^{22} \text{ cm}^{-2}$ (Sakano et

¹Visiting Astronomer, Kitt Peak National Observatory, National Optical Astronomy Observatories, which is operated by the Association of Universities for Research in Astronomy, Inc. (AURA) under cooperative agreement with the National Science Foundation.

²Visiting Astronomer, University of Hawaii Observatory.

³Research Fellow of the Japan Society for the Promotion of Science.

⁴Hubble Fellow.

al. 1998). We have found a galaxy with $R = 15.62$ mag near the center of the error circle for the X-ray source. The optical spectrum of the galaxy shows only narrow emission lines whose ratios correspond to those of a type 2 Seyfert galaxy at $z = 0.072$, implying an absorption-corrected X-ray luminosity of 2×10^{43} erg sec $^{-1}$ (2–10 keV) and $M_B = -20.93$ mag. A radio point source is also associated with the center of the galaxy. We thus identify the X-ray source with this galaxy as an obscured AGN. The hidden nature of the nucleus of the galaxy in the optical band is consistent with the X-ray spectrum. These results support the idea that the obscured AGNs/QSOs contribute significantly to the cosmic X-ray background in the hard band at the faint flux level.

Subject headings: galaxies: active — galaxies: individual (AX J131501+3141) — galaxies: Seyfert — X-rays: galaxies

1. Introduction

Many efforts have been made to understand the origin of the cosmic X-ray background (CXB). Recently in the *ROSAT* deep surveys, $\sim 60\%$ of the CXB in the 0.5–2 keV band has been resolved into discrete sources (Hasinger et al. 1993). On the other hand, in the harder 2–10 keV band, where the bulk of the energy density of the CXB resides, only $\sim 3\%$ of the CXB was resolved into discrete sources (Piccinotti et al. 1982). In the soft X-ray band (0.5–2 keV) type 1 AGNs are the main contributors to the CXB within a flux level of 1.6×10^{-15} erg cm $^{-2}$ s $^{-1}$ (McHardy et al. 1998). However, they have X-ray power-law spectra ($\Gamma = 1.7$, Turner & Pounds 1989) significantly softer than that of the CXB in the 2–10 keV band ($\Gamma = 1.4 \sim 1.5$, Gendreau et al. 1995, Ishisaki 1996). Therefore, there must be objects which have harder X-ray spectra than type 1 AGNs and contribute significantly to the CXB in this band.

To study the nature of X-ray sources in the hard band, we are now conducting an unbiased large and deep survey with the *ASCA* (Tanaka, Inoue, & Holt 1994) near the north Galactic pole. This is the Large Sky Survey (hereafter LSS, Inoue et al. 1996, Ueda 1996, Ueda et al. 1998). The flux limit of the LSS ($\sim 1 \times 10^{-13}$ erg cm $^{-2}$ s $^{-1}$ in the 2–10 keV band) is 100 times deeper than the *HEAO1* A2 survey, which was the deepest survey in the hard band (Piccinotti et al. 1982). We have surveyed 7.0 deg 2 and detected 43 sources above the 4σ level in the 2–10 keV band (Ueda 1996), corresponding to $\sim 30\%$ of the CXB in this band. The average photon index of the X-ray sources detected in the flux range between 1×10^{-13} and 5×10^{-13} erg cm $^{-2}$ s $^{-1}$ is $\Gamma = 1.5 \pm 0.2$ (Ueda et al. 1998), which is significantly harder than the spectra of X-ray sources detected in shallower surveys in the 2–10 keV band and close to that of the CXB. Thus, we are now revealing the presence of the hard sources which are responsible for most of the CXB energy density. Identification of these hard sources is clearly an important next step.

In the LSS, we have discovered the very hard X-ray source AX J131501+3141 whose power-law index is by far the hardest among the LSS sources. This object provides a good opportunity to examine the nature of the hard X-ray sources found in our systematic survey, and the result will be a key for the understanding of the origin of the CXB in the hard band. In this paper, we report on the optical identification and properties of AX J131501+3141, and their implications for the origins of the CXB. Results of the X-ray observations are presented in detail in a separate paper (Sakano et al. 1998). Throughout this paper, we use $q_0 = 0.5$ and $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2. Identification of AX J131501+3141

2.1. X-ray Properties

AX J131501+3141 is the hardest X-ray source detected in the LSS. The apparent X-ray spectrum from its initial observation can be fitted by a single power-law model with a photon index $\Gamma = -0.18 \pm 0.31$ ($f_E \propto E^{-(\Gamma-1)}$) in 0.7 – 10 keV band. Since the photon indices of the other X-ray sources detected in the LSS are distributed between 0.4 and 4.0 in the flux range from 1×10^{-13} to $1 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ (2–10 keV), the photon index of AX J131501+3141 is outstandingly hard.

Subsequently deeper X-ray observations of this object totaling 100,000 s were made to learn more details of its X-ray properties (Sakano et al. 1998). The results given in Sakano et al. (1998) are as follows. The flux of AX J131501+3141 in the 2–10 keV band is $\sim 5 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$, while in the 0.5 – 2 keV band no X-ray flux is detected above a 3 sigma level and the corresponding upper limit is $1.2 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$. The X-ray spectral shape seems to be a power law with a low energy cut-off at $\sim 2 - 3 \text{ keV}$; the best fitted parameters are $\Gamma = 1.5_{-0.6}^{+0.7}$ and absorbing column density $N_H = 6_{-2}^{+4} \times 10^{22} \text{ cm}^{-2}$ in the observed frame. The Galactic column density of neutral hydrogen in this direction is $N_{HI} = 1.1 \times 10^{20} \text{ cm}^{-2}$ (Stark et al. 1992). The deep observations were made at two epochs (1995 December and 1996 June) and 30% variability was detected. From the deep X-ray observations, the position of AX J131501+3141 is accurately determined to be $\alpha=13^h15^m00^s.9$, $\delta=31^\circ41'28''$ (J2000) with an error radius of $30''$ (95% confidence).

2.2. Optical Imaging Observations of AX J131501+3141

To identify the optical counterpart of AX J131501+3141, we made imaging observations with a Tektronix 2048×2048 CCD on the University of Hawaii 88" telescope in 1996 April. Images were obtained in two bands, *B* and *R*, with a spatial sampling of $0.''22 \text{ pix}^{-1}$. The exposure times were 20 and 15 min for the *B* and *R* bands, respectively. The FWHM of the seeing in the images was $1.''1$ in *B* and $1.''0$ in *R*.

Figure 1 shows the *R* band image centered on the position of AX J131501+3141 with its error circle of radius $30''$. In the error circle, there are one bright galaxy and a few faint objects. The

coordinates of the bright galaxy are $\alpha=13^{\text{h}}15^{\text{m}}01^{\text{s}}.15$, $\delta=31^{\circ}41'28''.1$ (J2000), which is only $3''.2$ from the best position of the X-ray source. Hence this is the most promising candidate for the optical counterpart. The galaxy has total magnitudes $R = 15.62 \pm 0.02$ and $B = 17.25 \pm 0.02$, and is cataloged as an S0 galaxy with $B = 17.2$ in Slezak et al. (1988). Hereafter, we call this bright galaxy “galaxy A”.

The other objects in the error circle are fainter than $B = 22.4$. Since type 1 AGNs are the most X-ray loud objects among the known extragalactic X-ray sources, we expect that the optical counterpart of AX J131501+3141 must be brighter than $B = 19.9$ on the basis of its 2–10 keV X-ray flux, assuming $\Gamma = 1.7$ together with the X-ray-to-optical flux ratio of type 1 AGNs identified in the Cambridge-Cambridge *ROSAT* Serendipity Survey (Boyle & di Matteo 1995). Since $B = 22.4$ is more than one order of magnitude fainter, none of these other objects are likely to be the optical counterpart.

The chance probability of one galaxy brighter than 16.0 mag falling in the error circle of the X-ray source is only 8×10^{-3} . This is obtained by adopting a surface density of such galaxies of 38 deg^{-2} (e.g., Stevenson, Shanks & Fong 1986). This low probability reinforces the conclusion that the galaxy A is the optical counterpart of AX J131501+3141.

2.3. A Radio Source in the Error Circle of AX J131501+3141

In the error circle of AX J131501+3141, a radio source is detected in the FIRST survey. The FIRST survey is a radio source survey conducted with the Very Large Array in the 1.4 GHz band with a 5σ limiting flux of 1 mJy (Becker, White, & Helfand 1995). The coordinates of the radio source are $\alpha=13^{\text{h}}15^{\text{m}}01^{\text{s}}.19$, $\delta=31^{\circ}41'29''.1$ (J2000), which is very close ($\sim 1''$) to the center of galaxy A (Table 1). The radio source is point-like and no structure can be seen. Considering the positional accuracy of the radio source ($\sim 1''$) and that of the optical galaxy center ($0''.8$), we regard the radio source as being associated with the center of galaxy A.

In the course of the optical follow-up program of the LSS, we examined the distribution of FIRST radio sources around LSS X-ray sources to evaluate the cross-correlation between radio and X-ray sources, and found a 3.5σ correlation within a radius of $0'.6$ from the X-ray sources (Akiyama et al. 1998). This radius corresponds to the error radius of the X-ray source positions. The presence of a significant correlation between the radio and X-ray sources implies that a FIRST radio source within $0'.6$ of an LSS X-ray source is the radio counterpart of the X-ray source with a probability of more than 80%. This is an additional reason why galaxy A, which is also the radio source detected near the center of the error circle of AX J131501+3141, is very likely to be the optical counterpart of the X-ray source.

In summary, considering all the results discussed in §2, we conclude that galaxy A and its central radio source are to be identified with AX J131501+3141. In the next section, we will present further evidence on the nature of galaxy A.

3. Nature of AX J131501+3141

3.1. Type 2 Seyfert at $z=0.072$

We made a spectroscopic observation of AX J131501+3141 (galaxy A) with the multi-slit spectrograph (CryoCam) on the KPNO Mayall 4m telescope on 1996 April 9. We used grism 730 and covered the spectral range from 6000 Å to 9000 Å with a resolution of 22 Å. We used a 2.''5 wide slit. The spectral sampling was 4.3 Å pix⁻¹ and the spatial resolution was 0.''84 pix⁻¹. The FWHM of the seeing was $\sim 1''$. The exposure time was 30 min. The spectrum was summed within a 4'' width along the slit direction (P.A. 62.°5) centered on the nucleus of the galaxy.

We detected emission lines of H α , [N II] $\lambda\lambda$ 6548,6583, and [SII] $\lambda\lambda$ 6717,6731 clearly but no broad H α emission line. The logarithmic line ratios are $\log([N II]\lambda 6583/H\alpha)=0.28$ and $\log([S II]\lambda 6717+6731/H\alpha)=-0.07$, which imply that this object is classified as a type 2 Seyfert or LINER (Veilleux & Osterbrock 1987). To be more definitive in the classification, spectrum in the blue wavelength region is needed. The redshift of this object turned out to be $z = 0.072$ and the resulting luminosities are $L_X = 2 \times 10^{43}$ ergs s⁻¹ in the 2–10 keV band after absorption-correction and $M_B = -20.93$ mag without K-correction. The power of the nuclear activity in the hard X-ray band is as large as that of a typical Seyfert galaxy (the knee of the luminosity function of type 1 AGNs is $L_X \sim 3 \times 10^{43}$ erg s⁻¹ in the 0.5–3.5 keV band according to Jones et al. 1997).

In order to obtain a spectrum in a blue region, we made another spectroscopic observation of galaxy A with the KPNO 2.1m telescope and the Goldcam spectrograph on 1997 June 8. Three 30 min exposures were taken through a 1.''9 slit at P.A.= 90°. The spectra covered the wavelength range 3866–7511Å with a spectral resolution of 5.1Å. The uncertainty in the wavelength scale is 0.1 Å. The flux from the nuclear region dominates the optical spectrum and it is only slightly resolved spatially in this observation. Therefore, we summed the spectrum using an optimal extraction method applied to a 13'' window along the slit. Atmospheric absorption band correction and flux calibration were then carried out with the help of spectra of standard stars. The resulting spectrum is shown in Figure 2. We can clearly see emission lines of [O II] λ 3727, H β , [O III] $\lambda\lambda$ 4959,5007, H α , [N II] $\lambda\lambda$ 6548,6583, and [SII] $\lambda\lambda$ 6717,6731.

The continuum contains a great deal of starlight, as evidenced by the Ca II H & K, H γ , Mg I b, and Na I D absorption lines. Also, the H β line shows an absorption plus emission structure. In order to obtain the more accurate measurements of the weak emission lines, we must subtract the stellar continuum spectrum. The model spectrum consists of an elliptical galaxy stellar template (NGC 5332) and a featureless power-law continuum, with the relative strengths of the two components adjusted to provide the smoothest fit to the continuum in the neighborhood of the emission lines. The spectrum obtained is also shown in Figure 2. Using the subtracted spectrum (Figure 2b dashed line), fluxes of the emission lines can be measured more accurately, though there are still obvious residuals due to the mismatch between the stellar populations of the two galaxies.

The resulting logarithmic line ratios turn out to be $\log([\text{N II}]\lambda 6583/\text{H}\alpha) = 0.06$, $\log([\text{S II}]\lambda 6717 + 6731/\text{H}\alpha) = -0.31$, and $\log([\text{O III}]\lambda 5007/\text{H}\beta) = 0.65$ (see Table 2). The ratio of $[\text{O III}]\lambda 5007/\text{H}\beta$ implies that the galaxy A is a Seyfert rather than LINER (Veilleux & Osterbrock 1987). These emission lines were resolved in the spectrum, with widths all approximately the same and consistent with $\text{FWHM} = 325 \pm 25 \text{ km s}^{-1}$. There is no hint of the presence of a broad-line component of $\text{H}\alpha$ or $\text{H}\beta$, even after subtraction of the continuum. Therefore we conclude galaxy A is classified as a type 2 Seyfert.

One interesting feature in the spectrum is that the intensity of $[\text{O II}]\lambda 3727$ emission line is strong in galaxy A. In the $[\text{O II}]\lambda 3727/[\text{O III}]\lambda 5007$ vs. $[\text{O III}]\lambda 5007/\text{H}\beta$ plane (Ferland & Netzer 1983), galaxy A is located between typical type 2 Seyferts and LINERs due to its large $\log([\text{O II}]\lambda 3727/[\text{O III}]\lambda 5007)$ values (0.004). After correcting for reddening using the large Balmer decrement ($\text{H}\alpha/\text{H}\beta = 7.58$), galaxy A falls among typical LINERs in this diagram because of the resulting larger value of $\log([\text{O II}]\lambda 3727/[\text{O III}]\lambda 5007) = 0.36$. There are two possible explanations for the ratio of the oxygen emission lines: 1) a small ionization parameter ($U \sim 10^{-3}$) in the nucleus, similar to LINERs, and 2) contamination by starforming regions in the disk of the galaxy, which would enhance $[\text{O II}]\lambda 3727$ relative to $[\text{O III}]\lambda 5007$. We favor the second explanation because a LINER classification is not supported by the other defining line ratio, $[\text{O I}]\lambda 6300/[\text{O III}]\lambda 5007$, which is less than $1/3$ in this spectrum, and also because the distribution of the $\text{H}\alpha$ emission along the slit in the spectra taken with the 2.1m telescope is slightly more extended than that of $[\text{N II}]\lambda 6583$. The spatial extent of the $\text{H}\alpha$ line is 4.2 pix FWHM ($3''.3$), while that of $[\text{N II}]\lambda 6583$ is 3.7 pix ($2''.9$). Unfortunately we cannot separate the nuclear component from an extended component in $[\text{O II}]\lambda 3727$ and $\text{H}\beta$ the signal-to-noise ratios of the lines are low and since the focus of the spectrum at the blue end is poor. The difference between $\text{H}\alpha$ and $[\text{N II}]\lambda 6583$ region is also confirmed in the spectrum taken with the 4m telescope. In addition, the color of the disk of galaxy A is $B - R = 1.3$, which is bluer than the bulge of the galaxy. These results suggest there is some contamination to the nuclear emission lines by star formation in the disk of the galaxy.

3.2. Hidden Nucleus of the Galaxy

The optical image (Figure 1) of AX J131501+3141 shows no point-like structure at the center of the galaxy, suggesting that the majority of the optical continuum photons in the central region of the galaxy originate in stars in the bulge. The column density to the nuclear X-ray source, $N_{\text{H}} = 6 \times 10^{22} \text{ cm}^{-2}$ (Sakano et al. 1998), corresponds to $A_V \approx 33 \text{ mag}$ according to the relations $N_{\text{HI}+\text{H}_2}/E(B - V) = 5.8 \times 10^{21} \text{ cm}^{-2} \text{ mag}^{-1}$ (Bohlin, Savage, & Drake 1978) and $A_V/E(B - V) = 3.14$ (Seaton 1979), which is large enough to hide the optical nuclear source. Thus the absence of a blue point-like nuclear source in the optical light is consistent with the large column density derived from the X-ray spectrum.

The large Balmer decrement ($\text{H}\alpha/\text{H}\beta = 7.58$) could be the result of extinction by a material distributed within or around the narrow-line region, rather than on the scale of the galaxy

(e.g., the disk of the edge-on galaxy). If the large Balmer decrement is the result of reddening, and we assume that the intrinsic ratio of $H\alpha$ to $H\beta$ is 3.1, the color excess $E(B - V)_{\text{Balmer}}$ is required to be 0.82. This value corresponds to the column density to the narrow-line region of $N_{\text{HI}+\text{H}_2} = 4.8 \times 10^{21} \text{ cm}^{-2}$. On the other hand, the $B - R$ color in a $5''$ aperture centered on galaxy A is $1.96 \pm 0.03 \text{ mag}$, which is redder than those of normal elliptical galaxies ($B - R = 1.47 \pm 0.05 \text{ mag}$, Buta et al. 1994, Buta and Williams 1995). If the intrinsic bulge color is the same as a typical color of an elliptical galaxy, the color excess of the bulge $E(B - V)_{\text{bulge}}$ is 0.28, if we adopt the relation $E(B - V) = E(B - R)/1.76$ (Seaton 1979). This value is smaller than that to the narrow-line region. Thus the difference between $E(B - V)_{\text{Balmer}}$ and $E(B - V)_{\text{bulge}}$ suggests that dust could be distributed on the scale of the narrow-line region, and smaller than that of the bulge.

3.3. Spectral Energy Distribution

Using the photometric results from radio to the X-ray wavelength, we can compare the spectral energy distribution (SED) of galaxy A with other type 2 Seyferts. Photometric results on the galaxy are summarized in Table 3. [In the infrared we use an upper limit from the *IRAS* Faint Source Catalog (Moshir et al. 1992) in the region containing galaxy A]. Figure 3 shows the SED of galaxy A as well as those of type 2 Seyferts taken from Mas-Hesse et al. (1995). The SEDs are normalized such that they have the same flux density as galaxy A at the optical wavelength ($\sim B$ band). Since the logarithmic flux density ratio between radio (1.4 GHz) and optical (B band) is -4.83 , which is similar to the other radio-quiet type 2 Seyferts, galaxy A is a radio-quiet object. The flux density in the hard X-ray band is located within a scatter of other type 2 Seyferts, galaxy A has a typical value of the flux density ratio of the hard X-ray to optical region as type 2 Seyferts.

3.4. An Interacting Galaxy?

In Figure 4, we show a wide image of of AX J131501+3141. Around AX J131501+3141, we can see two other bright galaxies of similar size, one (galaxy B) is located at $67''$ east, and the other (galaxy C) is located at $55''$ south-southeast of AX J131501+3141. In course of the multi-slit spectroscopy, we also observed these galaxies. The results are shown in Table 2. Since galaxy C has $z = 0.189$, its closeness to galaxy A is just a chance projection. Galaxy B has almost the same redshift (0.072) as galaxy A. The velocity difference between galaxy A and galaxy B is 90 km s^{-1} , and their projected separation is $67''$, which corresponds to 124 kpc at $z = 0.072$. The magnitudes of galaxy B are $B = 18.08$ and $R = 16.90$, and its morphology is that of a normal spiral galaxy without any clear distortion. However, the spectrum of galaxy B has emission lines which are H II-region like (Veilleux & Osterbrock 1987). The $B - R$ color of its disk is 1.0 mag which is bluer than the disk of galaxy A. Thus these results would indicate enhanced star-formation in the

disk of galaxy B. There is a possibility that the interaction is a driver of both the nuclear activity in galaxy A, and the enhanced star-formation in the disk of the galaxy B.

4. Implications for the Nature of Hard X-ray Sources

The hardest X-ray source detected in the continuous 7.0 deg² area of the LSS has been identified with a type 2 Seyfert having the column density to the nucleus of $N_{\text{H}} = 10^{22.8} \text{ cm}^{-2}$ in the local universe ($z \leq 0.1$). Surface density of such object is consistent with recent models of the origin of the CXB by e.g., Madau et al. (1994) and Comastri et al. (1995) as described below. They propose that type 2 AGNs are the hard X-ray sources which need to exist to account for the hardness of the spectrum of the CXB, because type 2 AGNs have harder X-ray spectra than type 1 AGNs due to the absorption of their soft X-ray photons. In the surveyed area, we have estimated the number density of type 2 AGNs which have the column density of $N_{\text{H}} = 10^{22-23} \text{ cm}^{-2}$ and X-ray flux larger than $5 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ in the 2–10 keV band i.e., the flux of AX J131501+3141. According to the model of Comastri et al. (1995), the expected number is 1.2. Observed surface number density is one (AX J131501+3141), if there are no more such objects in the LSS region. In fact, we cannot expect the presence of more such objects because of the following two reasons. 1) The AGNs with $N_{\text{H}} = 10^{22-23} \text{ cm}^{-2}$ in the local universe should have very hard X-ray spectra as like the hardest source; but no such hard source is found in the LSS region above $5 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ (Ueda 1996). 2) High redshift AGNs with $N_{\text{H}} = 10^{22-23} \text{ cm}^{-2}$ have softer X-ray spectra than the hardest source in the observed frame due to K-correction and may exist in the LSS region. However the probability that such high- z object exists is very low. Because the predicted redshift distribution of absorbed AGNs above $5 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ (2–10 keV) is peaked in $z \leq 0.1$ and declining with z in $z \geq 0.1$, if we adopt the evolution of X-ray luminosity function for type 1 AGNs (Jones et al. 1997) and assume the number ratio of type 1 to type 2 AGNs is constant with z .

In other surveys some hard X-ray sources have been also identified with absorbed AGNs. In Iwasawa et al. (1997), a hard X-ray source (AX J1749+684) detected serendipitously was identified with a nearby ($z = 0.05$) obscured AGN. The hard X-ray flux of the object is $9.6 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$, 2 times brighter than AX J131501+3141. A fainter hard X-ray source (AX J08494+4454) with $f_x = 9.0 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ in the 2–10 keV band, was identified with a type 2 QSO at $z = 0.9$ in course of optical follow-up observations of the *ASCA* Lynx deep survey (Ohta et al. 1996). A common feature of these three type 2 AGNs is that they are hard X-ray sources with radio detections. The α_{rx} ($f_\nu \propto \nu^{-\alpha}$) values of these three objects are 0.67 (AX J131501+3141), 0.67 (AX J1749+684) and 0.60 (AX J08494+4454). The agreement of these indices suggests that a hard X-ray survey together with a radio survey is a fairly efficient way to select obscured type 2 AGNs.

Recently, in some optical follow-up programs for the *ROSAT* deep fields, faint X-ray sources have been identified with narrow emission-line galaxies (NELGs) such as type 2 Seyferts,

star-forming galaxies, and LINERs (Boyle et al. 1995, Griffiths et al. 1996, McHardy et al. 1998). These objects are thought to be possible candidates for hard sources, but their X-ray spectra in the hard band have not been established. With respect to their soft X-rays, their 0.5–2 keV to optical flux ratios are one order of magnitude larger than those of normal galaxies and one order of magnitude smaller than those of QSOs (Griffiths et al. 1996). If we use the upper-limit of AX J131501+3141 in the 0.5–2 keV band, we find that its X-ray to optical flux ratio is more than 10 times smaller than those of typical NELGs, and closer to the normal galaxies ($\log[f_x(0.5 - 2keV)/f_B] = -1.52$). Thus AX J131501+3141 may be different from the NELGs detected in the *ROSAT* surveys. However, it is possible that we are seeing the same population of objects at lower redshifts. To establish the nature of the hard X-ray sources, identification of all the X-ray sources detected in the LSS is clearly necessary.

MA, KO, and TY would like to thank S.Okamura, M.Sekiguchi and the MOSAIC CCD camera team, and staff members of the KISO observatory for their support during the imaging observations. MA thanks K.Aoki and T.T.Takeuchi for their help. KO is grateful to the hospitality during his stay at the IfA, UH, where a part of this work was done. This research has made use of NASA/IPAC Extragalactic Database (NED), which is operated by the Jet Propulsion Laboratory, Caltech, under contract with the National Aeronautics and Space Administration. MA and MS acknowledge support from an Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists. The optical follow-up program is supported by grand-in-aids from the Ministry of Education, Science, Sports and Culture of Japan (06640351, 08740171, 09740173) and from the Sumitomo Foundation.

REFERENCES

- Akiyama, M., et al. 1998, in preparation
- Becker, R. H., White, R. L., & Helfand, D. J. 1995, *ApJ*, 450, 559
- Bohlin, R.C., Savage, B.D., & Drake, J.F. 1978, *ApJ*, 224, 132
- Boyle, B. J., & di Matteo, T. 1995, *MNRAS*, 277, L63
- Boyle, B. J., McMahon, R. G., Wilkes, B. J., & Elvis, M. 1995, *MNRAS*, 276, 315
- Buta, R., Mitra, S., deVaucouleurs, G., & Corwin Jr, H. G. 1994, *AJ*, 107, 118
- Buta, R., & Williams, K.L. 1995, *AJ*, 109, 543
- Comastri, A., Setti, G., Zamorani, G., & Hasinger, G. 1995, *A&A*, 296, 1
- Ferland, G. J., & Netzer, H. 1983, *ApJ*, 264, 105
- Gendreau, K. C., et al. 1995, *PASJ*, 47, L5
- Griffiths, R.E., Della Ceca, R., Georgantopoulos, I., Boyle, B.J., Stewart, G.C., Shanks, T., & Fruscione, A. 1996, *MNRAS*, 281, 71
- Hasinger, G., Burg, R., Giacconi, R., Hartner, G., Schmidt, M., Trümper, J., & Zamorani, G. 1993, *A&A*, 275, 1 [Erratum 1994, *A&A*, 291, 348]
- Inoue, H., Kii, T., Ogasaka, Y., Takahashi, T., Ueda, Y. 1996, in *Proc 'Röntgenstrahlung from the Universe'*, eds. Zimmermann, H.U., Trümper, J., and Yorke, H. MPE Report 263, 323
- Ishisaki, Y. 1996, PhD. thesis, University of Tokyo
- Iwasawa, K., Fabian, A.C., Brandt, W.N., Crawford, C.S., and Almaini, O., 1997, *MNRAS*, 291, L17
- Jones, L. R., et al. 1997, *MNRAS*, 285, 547
- Madau, P., Ghisellini, G., & Fabian, A.C., 1994, *MNRAS*, 270, L17
- Mas-Hesse, J. M., Rodriguez-Pascual, P. M., Sanz, L., Mirabel, I. F., Wamsteker, W., Makino, F., & Ohtani, C. 1995, *A&A*, 298, 22
- McHardy, I. M., et al. 1998, *MNRAS*, accepted
- Moshir, M., et al. 1992, *Explanatory Supplement to the IRAS Faint Source Survey, Version 2*, JPL D-10015 8/92 (Pasadena: JPL)
- Ohta, K., Yamada, T., Nakanishi, K., Ogasaka, Y., Kii, T., & Hayashida, K. 1996, *ApJ*, 458, L57

- Piccinotti, G., Mushotzky, R. F., Boldt, E. A., Holt, S. S., Marshall, F. E., Serlemitsos, P. J., & Shafer, R.A. 1982, *ApJ*, 253, 485
- Sakano, M., et al. 1998, *ApJ*, submitted
- Seaton, M. J. 1979, *MNRAS*, 187, 73p
- Slezak, E., Mars, G., Bijaoui, A., Balkowski, C., & Fontanelli, P. 1988, *A&AS*, 74, 83
- Stark, A. A., Gammie, C. F., Wilson, R. W., Bally, J., Linke, R. A., Heiles, C., and Hurwitz, M. 1992, *ApJS*, 79,77
- Stevenson, P. R. F., Shanks, T., & Fong, R. 1986, in *Spectral Evolution of Galaxies*, eds. C. Chiosi & A. Renzini (Reidel), 439
- Tanaka, Y., Inoue, H., & Holt, S. S. 1994, *PASJ*, 46, L37
- Turner, T. J., & Pounds, K. A. 1989, *MNRAS*, 240, 833
- Ueda, Y. 1996, PhD. thesis, University of Tokyo
- Ueda, Y., et al. 1998, *Nature*, accepted
- Veilleux, S., & Osterbrock, D. E. 1987, *ApJS*, 63, 295

Fig. 1.— R band image with the error circle of $30''$ radius centered on the X-ray position of AX J131501+3141. North is up and east is left.

Fig. 2.— a) Optical spectrum of the nuclear region of galaxy A. The abscissa shows observed wavelength in $\text{\AA}$ and the ordinates shows flux density in $10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$. Identified emission and absorption lines are marked at the top of the figure. b) Close up view of the optical spectra. The abscissa and ordinates are same as in a). Left panel shows $H\beta$ region, right panel shows $H\alpha$ region. Dashed line in each panel shows the spectra after subtracting of the model stellar continuum (see text).

Fig. 3.— The spectral energy distribution of AX J131501+3141. Large filled rectangles are data points of AX J131501+3141. Large filled triangle is the upper limit to the infrared emission. Other small marks represent SEDs of type 2 Seyferts from Mas-Hesse et al.(1995): open rectangles: Mrk 348, crosses: NGC 1068, filled rectangles: NGC 2110, open pentagons: Mrk 3, open hexagons: NGC 4388, filled hexagons: NGC 4507, open triangles: NGC 5506, arrows: NGC 5674, open circles: NGC 7582.

Fig. 4.— R band image around AX J131501+3141. Galaxies A, B, and C are marked. North is up and east is left.

band	name	coordinate (J2000)		uncertainty	distance ^a
X-ray	AX J131501+3141	13 ^h 15 ^m 00. ^s 9	31°41′28″	30″	3.″2
radio	FIRST source	13 ^h 15 ^m 01. ^s 19	31°41′29.″1	1″	1.″1
optical	galaxy A	13 ^h 15 ^m 01. ^s 15	31°41′28.″1	0.″8	—

Table 1: Coordinates of the X-ray, radio, and optical source.

^aDistance from the optical center of galaxy A.

No.	redshift	B^a	R^a	$\log([\text{NII}]/\text{H}\alpha)$	$\log([\text{SII}]/\text{H}\alpha)$	$\log([\text{OIII}]/\text{H}\beta)$
galaxy A	0.07203 ± 0.00009	17.25	15.62	0.06^e	-0.31^e	0.65^e
galaxy B	0.072	18.08	16.90	-0.39^f	-	–
galaxy C	0.189	18.75	17.03	0.0^f	-	–

Table 2: Results of the spectroscopic observations.

^aTotal magnitude.

^b $[\text{NII}]\lambda 6583$

^c $[\text{SII}]\lambda 6717+6731$

^d $[\text{OIII}]\lambda 5007$

^eLine ratio after subtraction of stellar continuum.

^fLine ratio from multi-slit spectroscopy without subtraction of stellar continuum.

name	band	frequency log(Hz)	observed flux density	$\log(\nu L_\nu)^a$
FIRST	1.4 GHz	9.15	3.83 mJy	39.08
<i>IRAS</i>	60 μ m	12.70	< 0.1 Jy	< 44.08
optical	<i>R</i> band	14.66	15.62 mag	44.21
	<i>B</i> band	14.83	17.25 mag	43.91
<i>ASCA</i>	2.0 keV	17.68	9.6×10^{-15b}	41.7
	10.0 keV	18.38	6.4×10^{-14b}	43.2

Table 3: Photometry of AX J131501+3141.

^aUnit in erg s⁻¹.

^bFlux density in erg s⁻¹ cm⁻² keV⁻¹ without absorption-correction.







